

# GÖDEL UNIVERSE AND SHEARS

Rainer Burghardt\*

Keywords: Gödel universe, shears, differential rotation law.

## Contents

1.	Introduction .....	2
2.	The locally non-rotating system .....	2
3.	Field equations for shears .....	4
4.	Conclusions.....	6
5.	References.....	7

It is shown that the Gödel universe is not rotating rigidly but has a differential rotation law. Locally non-rotating observers are subject to shears. The field strengths for the shears satisfy the field equations.

---

\* e-mail: [arg@aon.at](mailto:arg@aon.at), home page: <http://arg.or.at/>

# 1. INTRODUCTION

In a previous paper [1] we have shown that the Gödel metric can be re-interpreted by rescaling the variables. It has turned out that the rotation of the Gödel universe is not rigid, but has a differential rotation law. The two rotational distributions sum to a constant rotational field strength, usually thought to be proportional to a constant angular velocity. As a consequence of the assumption that a differential rotation law is valid it should be possible to choose a locally non-rotating reference system in which the observers are subject to shears. In the following we will show that the Einstein field equations provide these shears.

In Sec. 2 we introduce a locally non-rotating reference system for the Gödel metric, and in Sec. 3 we evaluate the field equations for the shears.

## 2. THE LOCALLY NON-ROTATING SYSTEM

Gödel [2] has given the metric of a rotating universe in the form

$$ds^2 = 4a^2 \left[ dr^2 + dy^2 + (\text{sh}^2 r - \text{sh}^4 r) d\phi^2 - 2\sqrt{2} \text{sh}^2 r d\phi dt - dt^2 \right], \quad (2.1)$$

satisfying the Einstein field equations for a rotating dust-filled universe. Rescaling and renaming the variables

$$a \rightarrow \mathcal{R}, \quad 2ay \rightarrow z, \quad 2at \rightarrow t, \quad 2r \rightarrow \eta, \quad r \rightarrow \chi$$

one can write this metric as

$$ds^2 = \mathcal{R}^2 d\eta^2 + dz^2 + \mathcal{R}^2 \text{sh}^2 \eta d\phi^2 - \left[ 2\sqrt{2} \mathcal{R} \text{sh}^2 \chi d\phi + dt \right]^2 \quad (2.2)$$

and with an additional substitution<sup>1</sup>

$$\text{th}\theta = \sqrt{2} \text{th}\chi \quad (2.3)$$

---

<sup>1</sup>  $\text{sh}^2\theta = \frac{2\text{sh}^2\chi}{1-\text{sh}^2\chi}$ ,  $\text{ch}^2\theta = \frac{\text{ch}^2\chi}{1-\text{sh}^2\chi}$ ,  $\text{th}^2\theta = 2\text{th}^2\chi$ ,  $\text{sh}\eta = 2\text{sh}\chi\text{ch}\chi$

$$ds^2 = R^2 d\eta^2 + dz^2 + R^2 \text{sh}^2 \eta d\phi^2 - [R \text{sh} \eta \text{th} \theta d\phi + dt]^2 \quad (\text{A}) . \quad (2.4)$$

We rearrange the metric as

$$ds^2 = R^2 d\eta^2 + dz^2 + \left[ 2R \text{sh} \chi \sqrt{1 - \text{sh}^2 \chi} d\phi - \frac{\sqrt{2} \text{sh} \chi}{\sqrt{1 - \text{sh}^2 \chi}} dt \right]^2 - \frac{ch^2 \chi}{1 - \text{sh}^2 \chi} dt^2 \quad (\text{B}) . \quad (2.5)$$

$$ds^2 = R^2 d\eta^2 + dz^2 + \left[ R \text{sh} \eta \frac{1}{\text{ch} \theta} d\phi - \text{sh} \theta dt \right]^2 - ch^2 \theta dt^2$$

The metric in the form (B) can be derived from (A) by a Lorentz transformation

$$L_2^{2'} = \text{cosi} \theta, \quad L_4^{2'} = \text{sini} \theta, \quad L_2^{4'} = -\text{sini} \theta, \quad L_4^{4'} = \text{cosi} \theta, \quad (2.6)$$

wherein  $\theta$  is the rapidity of the motion. The matter of the universe is in relative motion with respect to new observers. The velocity of the matter relative to these observers is

$$'u_m' = \{0, \text{sini} \theta, 0, \text{cosi} \theta\} = \{0, i\alpha \omega \sigma, 0, \alpha\} . \quad (2.7)$$

We have identified the transformation parameters with the physical quantities

$$\text{cosi} \theta = \text{ch} \theta = \alpha, \quad \text{sini} \theta = \text{sh} \theta = i\alpha \omega \sigma . \quad (2.8)$$

The circular velocity of the matter and the distance from the rotation axis are

$$\omega \sigma = \text{th} \theta, \quad \sigma = R \text{sh} \eta . \quad (2.9)$$

Evidently, the matter is rotating with the velocity of light ( $\text{th} \theta = 1$ ) at the *cut-off radius*

$$\sigma_c = \frac{1}{\omega_c} \quad (2.10)$$

where the rapidity is infinitely large. Beyond  $\sigma_c$  the motion is tachyonic and gives rise to acausalities and CTCs. Cooperstock [3] has mentioned that the light cones touch the world lines in that case. Pfarr [4] discussed closed null lines and paths with velocities larger than light.

The angular velocity

$$\omega(\chi) = \frac{\sqrt{2} \text{th} \chi}{R \text{sh} \eta} = \frac{1}{\sqrt{2} R \text{ch}^2 \chi} \quad (2.11)$$

depends on the distance to the rotation axis. Equ. (2.11) is the ansatz for the *differential rotation law*. In our previous paper [1] we have shown that the total rotational force has two contributions: a Coriolis-like one and a second, invoked by  $d\omega$ . Both contributions depend on the radial coordinate, but they sum to the total rotational force. This force is constant over all the space. This might pretend a constant rotational velocity.

Our aim is to show that in the local non-rotating system (system B) no Coriolis field strengths appear and that the observer experiences shears. We also show that the field strengths responsible for the shears, satisfy the field equations. Thus, we present a good argument for the assertion that the Gödel universe has to be described by a differential rotation law.

### 3. FIELD EQUATIONS FOR SHEARS

In paper [1] we have set up the field equations for the system (A). The evaluation of the field equations based on quantities deduced from the local non-rotating metric leads to a different structure. Rewriting the metric (B) once more with the physical quantities (2.9) we obtain

$$ds^2 = \mathcal{R}^2 d\eta^2 + dz^2 + \left[ \alpha^{-1} \sigma d\varphi + i \alpha \omega \sigma dx^4 \right]^2 + \alpha^2 dx^{4^2} . \quad (3.1)$$

In non-relativistic approximation the metric reads as

$$ds^2 = \mathcal{R}^2 d\eta^2 + dz^2 + \left[ \sigma (d\varphi - \omega dt) \right]^2 - dt^2 . \quad (3.2)$$

With the 4-bein vectors

$$\begin{aligned} \mathbf{e}_1^1 = 1, \quad \mathbf{e}_2^2 = \frac{\sigma}{\alpha}, \quad \mathbf{e}_3^3 = 1, \quad \mathbf{e}_4^4 = i \alpha \omega \sigma, \quad \mathbf{e}_4^4 = \alpha \\ \mathbf{e}_1^1 = 1, \quad \mathbf{e}_2^2 = \frac{\alpha}{\sigma}, \quad \mathbf{e}_3^3 = 1, \quad \mathbf{e}_4^2 = -i \alpha \omega, \quad \mathbf{e}_4^4 = \frac{1}{\alpha} \end{aligned} \quad (3.3)$$

we can calculate the quantities

$$\begin{aligned} A_{2m}^2 = \bar{B}_m = B_m - F_m - D_m, \quad A_{4m}^4 = G_m = F_m + D_m \\ D_{mns} = 2 \left[ D_{(mn)} u_s - D_{(ms)} u_n + D_{[ns]} u_m \right] \\ F_m = \alpha^2 \omega^2 \sigma \sigma_{|m}, \quad D_m = \alpha^2 \omega \omega_{|m} \sigma^2 \end{aligned} \quad (3.4)$$

These quantities have only a few non-vanishing components

$$B_1 = \frac{1}{R} \text{cth}\eta, \quad F_1 + D_1 = \omega \text{sh}\theta \text{ch}\theta, \quad D_{12} = i\omega [\text{sh}^2\theta - \text{sh}^2\chi], \quad D_{21} = 0. \quad (3.5)$$

$\bar{B}_m$  is the curvature quantity of the 3-dimensional geometry,  $F_m$  the centripetal force and  $D_m$  a contribution from the differential rotation law. The  $D_{mn}$  are the shears. The Ricci-rotation coefficients are composed of all these quantities

$$A_{mn}{}^s = b_m \bar{B}_n b^s - b_m b_n \bar{B}^s + D_{mn}{}^s + u_m G_n u^s - u_m u_n G^s$$

$$b_m = \{0, 1, 0, 0\}, \quad u_m = \{0, 0, 0, 1\} \quad (3.6)$$

The  $u_m$  are the 4-velocities of the observers of the *locally non-rotating reference system* (LNR). With these connexion coefficients one has

$$u_{m||n} u^n = -(F_m + D_m), \quad u_{[\alpha||\beta]} = 0, \quad u_{(\alpha||\beta)} = -2D_{(\alpha\beta)}. \quad (3.7)$$

The last two relations show that the LNR observers experience no Coriolis-like forces but shears.

Defining the graded covariant derivatives

$$\bar{B}_{n||m} = \bar{B}_{n|m}, \quad G_{n||m} = G_{n|m} - \bar{B}_{mn}{}^s G_s, \quad (3.8)$$

we can write the Ricci as

$$R_{mn} = - \left[ \bar{B}_{n||m} + \bar{B}_n \bar{B}_m \right] - \left[ G_{n||m} + G_n G_m + 2D_n{}^s D_{ms} \right]$$

$$- b_m b_n \left[ \left( B_{||s}^s + B^s B_s \right) - \left( G_{||s}^s + G^s G_s + 2D^{sr} D_{sr} \right) \right]$$

$$- u_m u_n \left[ G_{||s}^s + G^s G_s + 2D^{sr} D_{sr} \right]$$

$$- 2u_{(m} \left[ D_{n)||s}^s + D^s_{(n} \bar{B}_{s)} \right] \quad (3.9)$$

Gödel's matter tensor is transformed with (2.6) to

$$T_{mn} = \mu_0 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -\alpha^2 \omega^2 \sigma^2 & 0 & -i\alpha^2 \omega \sigma \\ 0 & 0 & 0 & 0 \\ 0 & -i\alpha^2 \omega \sigma & 0 & \alpha^2 \end{pmatrix} = \mu_0 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -\text{sh}^2\theta & 0 & -i\text{sh}\theta \text{ch}\theta \\ 0 & 0 & 0 & 0 \\ 0 & -i\text{sh}\theta \text{ch}\theta & 0 & \text{ch}^2\theta \end{pmatrix}. \quad (3.10)$$

Gödel used the extended field equations with the cosmological constant

$$R_{mn} - \frac{1}{2} R g_{mn} - \lambda g_{mn} = -\kappa T_{mn} . \quad (3.11)$$

As  $\lambda = -\frac{R}{2}$  one is left with

$$R_{mn} = -\kappa T_{mn} . \quad (3.12)$$

For the radial part of the field equations one obtains

$$G^s_{||s} = \kappa \mu_0 \alpha^2 - G^s G_s - 2D^{sr} D_{sr} \quad (3.13)$$

and for the shears

$$D^s_{n||s} = -\kappa \mu_0 i \alpha^2 \omega \sigma b_n - D^s_n \bar{B}_s . \quad (3.14)$$

The radial field strengths are coupled to the matter density and to the field energy including the shears. The shears have the matter current and a field contribution as source. Finally, we add some references concerning Gödel's model [5 – 57].

## 4. CONCLUSIONS

We have shown that our assertion that an observer in the Gödel universe can experience shears is aided by the introduction of a locally non-rotating system. If the metric is reordered and disassembled in 4-bein fields in a proper way one obtains from the Ricci-rotations coefficients field strengths which are responsible for the shears of the observer fields. These shears are closely related to a differential rotation law and satisfy the field equations.

## 5. REFERENCES

1. Burghardt, R., *Constructing the Gödel universe*. <http://arg.or.at/Wpdf/WGoe.pdf>
2. Gödel, K., *An example of a new type of cosmological solutions of Einstein's field equations of gravitation*. *Rev. Mod. Phys.* **21**, 447, 1949
3. Cooperstock F. I., Tieu S., *Closed time-like curves re-examined*. gr-qc/0405114
4. Pfarr J.; *Time travel in Gödel's space*. *GRG* **13**,1073, 1981
5. Bampi F., Zordan C., *A note on Gödel's metric*. *GRG* **9**, 393, 1978
6. Banjeree A., Banjeri S., *Stationary distributions of dust and electromagnetic fields in general relativity*. *Proc. Phys. Soc.* **1**, 118, 1968
7. Barrow J. D., Dąbrowski M. P., *Gödel universes in string theory*. gr-qc/9803048
8. Barrow J. D., *Dynamics and stability of the Gödel universe*. *Class. Quant. Grav.* **21**, 1773, 2004
9. Bonnor W. B., Santos N. O., McCallum M. A. H., *An exterior for the Gödel spacetime*. gr-qc/9711011
10. Bonnor W. B., *Closed timelike curves in general relativity*. gr-qc/0211051
11. Bray M., *Sur quelques univers magnétohydrodynamiques du type de Gödel*. *Compt. Rend.* **A 274**, 874, 1972
12. Bray M., *Quelques univers magnétohydrodynamiques du type de Gödel*. *Compt. Rend.* **A 274**, 809, 1972
13. Caldarelli M. M., Klemm D., *Supersymmetric Gödel-type universe in four dimensions*. hep-th/0310081
14. Chicone C., Mashhoon B., *Explicit Fermi coordinates and tidal dynamics in de Sitter and Gödel spacetime*. gr-qc/0511129
15. Chandrasekhar S., Wright J. P., *The geodesics in Gödel's universe*. *Proc. Nat. Ac. Sci.* **47**, 341, 1961
16. Clifton T., Barrow J. D., *The existence of Gödel, Einstein and de Sitter universes*. gr-qc/0511076
17. Dąbrowski M. P., Garecki J., *Energy-momentum and angular momentum of Gödel universes*. gr-qc/0309064
18. Das S., Gegenberg J., *Gravitational non-commutativity and Gödel-like spacetimes*. gr-qc/0407053
19. Drukker N., Fiol B., Simon J., *Gödel's universe in a supertube shroud*. gr-qc/0306057
20. Drukker N., Fiol B., Simon J., *Gödel type universe and the Landau problem*. hep-th/0309199
21. Dryuma V., *On the Riemannian extension of the Gödel space time metric*. gr-qc/0511165
22. Figueiredo B. D. B., *Gödel type spacetimes and the motion of charged particles*. *Class. Quant. Grav.* **15**, 3849, 1998
23. Fonseca-Neto J. B., Romero C., Dahia F., *Embedding Gödel's universe in five dimensions*. gr-qc/0503122
24. Gürses M., Karasu A., Sarıoğlu Ö., *Gödel type of metrics in various dimensions*. hep-th/0312290
25. Kanti P., Vayonakis C. E., *Gödel-type universes in string-inspired charged gravity*. gr-qc/9905032
26. Kundt W., *Trägheitsbahnen in einem von Gödel angegebenen kosmologischen Modell*. *Z. f. Phys.* **145**, 611, 1956
27. Laurent B.E., Rosquist K., Sviestins E., *The behavior of null geodesics in a class of rotating space-time homogeneous cosmologies*. *GRG* **13**, 11, 1981
28. Novello M., Rebouças M. J., *The stability of a rotating universe*. *Astrophys. Journ.* **225**, 719, 1978
29. Novello M., Soares I. D., Tiomno J., *Geodesic motion and confinement in Gödel's universe*. *Phys. Rev.* **D 27**, 779, 1983
30. Maitra S. C., *Stationary dust-filled cosmological solution with  $\Lambda=0$  and without closed timelike lines*. *Journ. Math. Phys.* **7**, 1025, 1966
31. Ozsvath I., *Dust-filled universes of class II and class III*. *Journ. Math. Phys.* **11**, 2871, 1970
32. Ozsvath I., *Spatially homogeneous world models*. *Journ. Math. Phys.* **11**, 2860, 1970
33. Ozsvath I., *On spatially homogeneous rotating world models*. Preprint
34. Ozsvath I., *Lösungen der Einsteinschen Feldgleichungen mit einfach transitiver Bewegungsgruppe*. *Ak. Wiss. Lit. Mainz Nat. Kl.* **13**, 1002, 1962
35. Ozsvath I., Schücking E. L., *The finite rotating universe*. *Ann Phys.* **55**, 166, 1969
36. Panov V. F., *A cosmological model with rotation in the relativistic theory of gravity*. *Teor. Mat. Fiz.* **114**, 249, 1998
37. Paiva F. M., Rebouças M. J., Teixeira A. F. F., *Time travel in homogeneous Som-Raychaudhuri universe*. CBPF-NF-056/87

38. Patel L.K., Trivedi V. M., *Einstein-Gödel universe with an electromagnetic field.* Curr. Sci. Ind. **47**, 800, 1978
39. Radu E. *On a Gödel-type Euclidean solution.* GRG **31**, 287, 1999
40. Raval H. M., Vaidya P. C., *On a Gödel universe filled with charged incoherent matter.* Curr. Sci. Ind. **36**, 7, 1967
41. Raychaudhuri A.,K.,Thakurta S. N. G., *Homogeneous space times of Gödel type.* Phys. Rev. **D 22**, 802, 1980
42. Rebouças M. J., A rotating universe with violation of causality. Phys. Lett. 70 a, 161, 1979
43. Rebouças M. J., Aman J. E., Teixeira A. F. F., *A note on Gödel-type space-times.* Journ. Math. Phys. **27**, 1370, 1986
44. Rebouças M. J., Aman J. E., *Computer aided study of a class of Riemannian space-times.* CBPF-NF-041/86
45. Rebouças M. J., Tiomno J., *Homogeneity of Riemannian space-times of Gödel type.* Am. Phys. Soc. **28**, 1251, 1983
46. Romano A. E., *Derivation of Gödel type metric with isometry group  $SO(2,1) \times SO(2) \times R$ .* gr-qc/0208093 vs1
47. Romano A. E., Goebel C., *Gödel-type space-time metrics.* gr-qc/0208093 vs 3
48. Rooman M., Spindel Ph., *Gödel metric as a squashed anti-de Sitter geometry.* Class. Quant. Grav. **15**, 3241, 1998
49. Rosa V. M., Letelier P. S., *Stability of closed timelike curves in Gödel universe.* gr-qc/0703100
50. Sahdev D., Sundararaman R., Modgil M., S.,*The Gödel universe: A practical travel guide.* gr-qc/0611093
51. Sharif M., *Energy momentum in spacetime homogeneous Gödel-type metrics.* gr-qc/0310018]
52. Sharif M., *Energy and momentum in spacetime associated with Gödel universe.* Int. Journ. Mod. Phys. **18**, 4361, 2003
53. Silk J., *Local irregularities in a Gödel universe.* Astr. Phys. Journ. **143**, 689, 1966
54. Stein H., *On the paradoxical time-structures of Gödel.* Phil. Sci. **11**, 21, 1977
55. Teixeira A. F. F., Rebouças M. J., Aman J. E., *Isometries of homogeneous Gödel-type spacetimes.* Phys. Rev. **D 32**, 3309, 1985
56. Vaidya P. C., *An Einstein-Gödel universe.* GRG **9**, 801, 1978
57. Wright J. P., *Solution of Einstein's field equations for a rotating, stationary and dust-filled universe.* Journ. Math. Phys. **6**, 103, 1965